

Literatur

- Anderson HV et al. Coronary artery flow velocity is related to lumen area and regional left ventricular mass. *Circulation* 2000; 102: 48–54.
- Baker KM, Booz GW, Dostal DE. Cardiac actions of angiotensin II: role of an intracardiac rennin-angiotensin system. *Annu Rev Physiol* 1992; 54: 227–241.
- Bhatia RS et al. Outcome of heart failure with preserved ejection fraction in a population-based study. *N Engl J Med* 2006; 355: 260–269.
- Brilla CG, Janicki JS, Weber KT. Impaired diastolic function and coronary reserve in genetic hypertension: role of interstitial fibrosis and medial thickening of intramyocardial coronary arteries. *Circ Res* 1991; 69: 107–115.
- Brilla CG, Funck RC, Rupp H. Lisinopril-mediated regression of myocardial fibrosis in patients with hypertensive heart disease. *Circulation* 2000; 102: 1388–1393.
- Ciaroni S, Cuenoud L, Bloch A. Clinical study to investigate the predictive parameters for the onset of atrial fibrillation in patients with essential hypertension. *Am Heart J* 2000; 139: 814–819.
- Drazner MH et al: Left ventricular hypertrophy is more prevalent in blacks than whites in the general population. The Dallas Heart Study. *Hypertension* 2005; 46: 124.
- Ezzati M et al; Comparative Risk Assessment Collaborating Group. Selected major risk factors and global and regional burden of disease. *Lancet* 2002; 360: 1347–1360.
- Frank D et al. Gene expression pattern in biomechanically stretched cardiomyocytes: Evidence for a stretch-specific gene program. *Hypertension* 2008; 51: 309–318.
- Frohlich ED et al. The heart in hypertension. *N Engl J Med* 1992; 327: 998–1008.
- Gaasch WH et al. Stress-shortening relations and myocardial blood-flow in compensated and failing canine hearts with pressure-overload hypertrophy. *Circulation* 1989; 79: 872–883.
- Galderisi M et al. Impact of ambulatory blood pressure on left ventricular diastolic dysfunction in arterial systemic hypertension. *Am J Cardiol* 1996; 77: 579–601.
- Galderisi M et al. Coronary flow reserve and myocardial diastolic dysfunction in arterial hypertension. *Am J Cardiol* 2002; 90: 860–864.
- Ghali JK et al. Impact of left ventricular hypertrophy on ventricular arrhythmias in the absence of coronary artery disease. *J Am Coll Cardiol* 1991; 17: 1277–1282.
- Hennersdorf MG, Strauer BE. Arterial hypertension and cardiac arrhythmias. *J Hypertens* 2001; 2: 167–177.
- Hittinger L et al. Hemodynamic mechanisms responsible for reduced subendocardial coronary reserve in dogs with severe left ventricular hypertrophy. *Circulation* 1995; 92: 978–986.
- Hunter WC. Role of myofilaments and calcium handling in left ventricular relaxation. *Cardiol Clin* 2000; 18: 443–457.
- Inouye I et al. Abnormal left ventricular filling: an early finding in mild to moderate systemic hypertension. *Am J Cardiol* 1984; 53: 120–126.
- Kannel WB, Schatzkin A. Sudden death: Lessons from subsets in population studies. *J Am Coll Cardiol* 1985; 5: 141–149.
- Kannel WB et al. Prevalence, incidence, prognosis and predisposing conditions for atrial fibrillation: population-based estimates. *Am J Cardiol* 1998; 82: 2–9.
- Lamb HJ et al: Diastolic dysfunction in hypertensive heart disease is associated with altered myocardial metabolism. *Circulation* 1999; 99: 2261–2267.
- Leite-Moreira AF, Gillebert TC. The physiology of left ventricular pressure fall. *Rev Port Cardiol* 2000; 19: 1015–1021.
- McLenachan JM et al. Ventricular arrhythmias in patients with hypertensive left ventricular hypertrophy. *N Engl J Med* 1987; 317: 787–792.
- Levy D et al. The progression from hypertension to congestive heart failure. *JAMA* 1996; 275: 1557–1562.
- Levy D et al. Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med* 1990; 322: 1561–1566.
- Messerli FH. Hypertension and sudden cardiac death. *Am J Hypertens* 1999; 12: 181S–188S.
- Nakao K et al. Myosin heavy chain gene expression in human heart failure. *J Clin Invest* 1997; 100: 2362–2370.
- Okin PM et al. Electrocardiographic strain pattern and prediction of new-onset congestive heart failure in hypertensive patients: The Losartan Intervention for Endpoint Reduction in Hypertension (LIFE) study. *Circulation* 2006; 113: 67.
- Owan TE et al. Trends in prevalence and outcome of heart failure with preserved ejection fraction. *N Engl J Med* 2006; 355: 251–259.
- Rockman HA, Koch WJ, Lefkowitz RJ. Seven transmembrane spanning receptors and heart function. *Nature* 2002; 415: 206–212.
- Rossi MA. Pathologic fibrosis and connective tissue matrix in left ventricular hypertrophy due to chronic arterial hypertension in humans. *J Hypertens* 1998; 16: 1031–1041.
- Sadoshima J, Izumo S. The cellular and molecular response of cardiac myocytes to mechanical stress. *Annu Rev Physiol* 1997; 59: 551–571.
- Schafer S et al. Left ventricular remodelling impairs coronary flow reserve in hypertensive patients. *J Hypertens* 2002; 20: 1431–1437.
- Schartzkopff B et al. Repair of coronary arterioles after treatment with perinopril in hypertensive heart disease. *Hypertension* 2000; 36: 220–225.
- Treasure CB et al. Hypertension and left ventricular hypertrophy are associated with impaired endothelium-mediated relaxation in human coronary resistance vessels. *Circulation* 1993; 87: 86–93.
- Unger T. The role of the renin-angiotensin system in the development of cardiovascular disease. *Am J Cardiol* 2002; 89: 3–10.
- Weber KT. Extracellular matrix remodelling in heart failure: a role for de novo angiotensin II generation. *Circulation* 1997; 96: 4065–4082.
- Weber KT. Targeting pathologic remodelling: concepts of cardioprotection and reparation. *Circulation* 2000; 102: 1342–1345.
- Youn HJ et al. Relation between flow reserve capacity of penetrating intramyocardial coronary arteries and myocardial fibrosis in hypertension: study using transthoracic Doppler echocardiography. *J Am Soc Echocardiogr* 2006; 19: 373–378.
- Yusuf S et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364: 937–952.
- Sipahi I et al. Effects of normal, pre-hypertensive, and hypertensive blood pressure levels on progression of coronary atherosclerosis. *J Am Coll Cardiol* 2006; 48: 833–838.